

Renormalization of the meson-baryon SU(3) Lagrangian in chiral perturbation theory to order q^3

GUIDO MÜLLER ^a

*Institut für Theoretische Kernphysik
Nussallee 14 - 16
53115 Bonn, Germany*

We perform the complete regularization of all Green functions with a single incoming and outgoing baryon to order q^3 in the low energy expansion for the SU(3) meson-baryon system in chiral perturbation theory. The divergences can be extracted in a chiral invariant manner by making use of the heat kernel representation of the propagators in d-dimensional Euclidean space.

In the arena of low energy physics it is much more useful to introduce an effective Lagrangian, which is written in terms of experimental degrees of freedom –mesons and baryons– and which encodes the symmetries of the underlying QCD interaction – specifically for our purpose chiral symmetry, which exists in the limit in which the quark mass can be taken as vanishing. Since we are interested in processes where the momenta are small, we can expand the Green functions in powers of the external fields. The low energy expansion involves two small parameters, the external momenta q and the quark masses $\mathcal{M}$. One expands in powers of these with the ratio $\mathcal{M}/q^2$ fixed. The low energy expansion is now obtained from a perturbative expansion of the meson EFT,

$$\mathcal{L}_M = \mathcal{L}_M^{(2)} + \mathcal{L}_M^{(4)} + \dots \quad (1)$$

where the index ($d = 2, 4, \dots$) denotes the low energy dimension. The leading term in the low energy expansion (1) can be easily written down in terms of the pseudoscalar Goldstone fields ($\phi = \pi, K, \eta$) which are collected in a 3×3 unimodular, unitary matrix $U(x)$,

$$U(\phi) = u^2(\phi) = \exp\{i\phi/F_0\} \quad (2)$$

transforming linearly under chiral symmetry. The lowest order Lagrangian consistent with Lorentz invariance, chiral symmetry, parity, G-parity and charge conjugation reads

$$\mathcal{L}_M^{(2)} = \frac{F_0^2}{4} (\langle \partial_\mu U \partial^\mu U^\dagger \rangle + 2B_0 \langle \mathcal{M}(U + U^\dagger) \rangle) \quad (3)$$

^ae-mail: mueller@pythia.itkp.uni-bonn.de

The scale F_0 can be identified with the pseudoscalar decay constant in the chiral limit and the constant B_0 is related to the quark condensate. One can now calculate tree diagrams using the effective Lagrangian $\mathcal{L}_M$ and derive all current algebra predictions. One has to include higher order corrections and meson loops, because tree diagrams are always real and thus unitarity is violated. Weinberg made the important observation that diagrams with L ($L = 1, 2, \dots$) meson loops are suppressed by powers of $(q^2)^L$ with respect to the leading term. In the meson sector we have a one-to-one correspondence between the number L of loops and the chiral dimension

$$D = 2 + \sum_d N_d(d-2) + 2L \quad (4)$$

where N_d denotes the number of vertices of dimension d . For a given S -matrix element, the chiral dimension D increases with the number of meson loops. It should be not surprising that a generic one-loop diagram in CHPT is divergent. The divergences can be taken care of once and for all by calculating the divergent part of the one-loop functional. Since it must be a local action with the symmetries of $\mathcal{L}_M^{(2)}$ and since it has chiral dimension $D = 4$, the corresponding Lagrangian must be of the general form as discussed in ref⁴.

$$\mathcal{L}_4 = \sum_{i=1}^{10} L_i P_i + \sum_{i=1}^2 H_i \tilde{P}_i \quad (5)$$

The renormalization procedure at one-loop level is then achieved by simply dividing the coefficients in (5) into a scale dependent finite and divergent part:

$$L_i = L_i^r(\mu) + \Gamma_i \Lambda. \quad (6)$$

The constants Γ_i serve to renormalize the infinities of the meson loops and the remaining finite pieces $L_i^r(\mu)$ have to be fixed phenomenologically or to be estimated by some model. With dimensional regularization, the divergent factor Λ is given by

$$\Lambda = \frac{\mu^{d-4}}{(4\pi)^2} \left\{ \frac{1}{d-4} - \frac{1}{2} [\log(4\pi) + 1 + \Gamma'(1)] \right\} \quad (7)$$

depending on the arbitrary renormalization scale μ . The sum of one-loop amplitudes and of tree amplitudes based on (5) is not only finite, but also independent of the scale μ . To evaluate these loop graphs one considers the neighbourhood of the solution to the classical equations of motion. The contribution to the one-loop graphs to the generating functional is given by

$$\exp\{iZ_{1-loop}\} = N[\det(dd + \sigma)]^{-\frac{1}{2}} \quad (8)$$

where the explicit definitions are given in ⁴. The determinant is UV divergent and produces poles at $d = 0, 2, 4, \dots$ in dimensional regularization and can be regularized via the heat kernel expansion.

Let us now consider the meson-baryon interaction in CHPT. Because of the different Lorentz structure of meson and baryon fields, the chiral expansion of $\mathcal{L}_{MB}$ contains terms of $O(q^n)$ for each positive integer n , unlike in the case of $\mathcal{L}_M$ where the chiral expansion proceeds in steps of two powers of q . So we have

$$\mathcal{L}_{MB} = \mathcal{L}_{MB}^{(1)} + \mathcal{L}_{MB}^{(2)} + \mathcal{L}_{MB}^{(3)} + \dots \quad (9)$$

The effective meson-baryon Lagrangian for processes with one incoming and one outgoing baryon is bilinear in the baryon field and starts with terms of dimension one,

$$\mathcal{L}_{MB}^{(1)} = \langle \bar{B} (i\not{\nabla} - m) B \rangle + \frac{D}{2} \langle \bar{B} \{ \not{\psi} \gamma_5, B \} \rangle + \frac{F}{2} \langle \bar{B} [\not{\psi} \gamma_5, B] \rangle \quad (10)$$

We collect the spin- $\frac{1}{2}$ baryon octet in the 3×3 matrix B . The lowest order meson-baryon Lagrangian contains two axial-vector coupling constants, denoted by D and F and the average baryon octet mass m in the chiral limit. In contrast, baryon propagators introduce the baryon mass as an additional dimensional quantity, which remains finite in the chiral limit. Consequently, we have not a correspondence between loop expansion and chiral dimension as in the meson sector expressed in eq.(4). With inspiration from heavy quark effective theory, Jenkins and Manohar ⁷ have reformulated baryon CHPT in precisely such a way as to transfer the baryon mass from the propagator to the vertices by defining velocity-dependent fields. The chiral dimension D for processes with exactly one baryon line running through the pertinent Feynman diagrams is given by ⁹

$$D = 2L + 1 + \sum_{d=2,4,6,\dots} (d-2) N_d^M + \sum_{d=1,2,3,\dots} (d-1) N_d^{MB} \geq 2L + 1 \quad (11)$$

with N_d^M (N_d^{MB}) counts the number of mesonic (meson-baryon) vertices of dimension d . In contrast to the relativistic formulation, chiral power counting in the heavy baryon formalism is completely analogous to the mesonic case. As for the lowest-order constants F_0, B_0 in (3), the coupling constants in $\mathcal{L}_{MB}^{(1)} + \mathcal{L}_{MB}^{(2)}$ are not renormalized in any order of the loop expansion, since $D \geq 3$. As in the meson-baryon sector we expand around the classical solution of motion and the corresponding one-loop generating functional consists of four pieces corresponding to two irreducible and two reducible diagrams ^{2 6 8}. It can be shown that the sum of the reducible diagrams is finite and scale

independent. The divergences of the one-loop graphs are therefore contained in the irreducible diagrams which are given by^{2,8}

$$Z_{\text{irr}}[j, R_v^a] = \int d^4x d^4x' d^4y d^4y' \bar{R}_v^a(x) S_{(1)}^{bc, \text{cl}}(x, y) \times \\ [\Sigma_2^{cd}(y, y) \delta(y - y') + \Sigma_1^{cd}(y, y')] S_{(1)}^{de, \text{cl}}(y', x') R_v^e(x') \quad (12)$$

Σ_1 corresponds to self-energy diagrams and Σ_2 contains tadpole diagrams. Note that the propagators and the vertices have the full tree-level structure attached to them as functionals of the external fields. The regularization in the meson-baryon sector is extremely tedious, because we are not able to express the one-loop functional in terms of a determinant unlike the meson sector. Nevertheless the divergences can be extracted in a chiral invariant manner by making use of the heat kernel representation of the propagators in d-dimensional Euclidean space. In the case of the self-energy contribution the divergences are due to the singular behaviour of the product of the meson and the baryon propagators in the coincidence limit. A non-trivial part of the regularization consists in finding a heat kernel representation for the full tree-level baryon propagator². The main difference between the SU(2) and the SU(3) calculation lies in the fact that the nucleons are in the fundamental representation of SU(2), while the baryons are in the adjoint representation of SU(3). This group theoretical difference in flavor space does not change the singular behaviour in coordinate space which leads to the divergences. The complete renormalization of Green functions for the pion-nucleon SU(2) interaction was done by Ecker² and we performed the regularization in SU(3) CHPT⁸. To separate the finite parts in dimensional regularization, we follow the conventions of⁵ to decompose the irreducible one-loop functional into a finite and a divergent part. Both depend on the scale μ . The generating functional can then be renormalized by introducing the counterterm Lagrangian⁸

$$\mathcal{L}_{MB}^{(3)\text{ct}}(x) = \frac{1}{(4\pi F_0)^2} \sum_{i=1}^{102} d_i \bar{H}_v^{ab}(x) \tilde{O}_i^{bc}(x) H_v^{ca}(x) \quad (13)$$

where the d_i are dimensionless coupling constants and the field monomials $\tilde{O}_i^{bc}(x)$ are of order q^3 . The low-energy constants d_i are decomposed in analogy to Eq.(6). The β_i are dimensionless functions of F and D constructed such that they cancel the divergences of the one-loop functional. The counterterms constitute a complete set for the renormalization of the irreducible tadpole and self-energy functional for *off-shell* baryons. These are the terms where the covariant derivative acts on the baryon fields. As long as one is only interested in Green functions with *on-shell* baryons, the number of terms can be reduced

considerably by invoking the baryon equation of motions. A further reduction in the number of terms could be achieved by use of the Cayley–Hamilton relation. Also, many of the terms given in the table refer to processes with at least three Goldstone bosons. These are only relevant in multiple pion or kaon production by photons or pions off nucleons¹⁰. The renormalized LECs $d_i^r(\mu)$ are measurable quantities. They satisfy the renormalization group equation. Therefore, the choice of another scale μ_0 leads to modified values of the renormalized LECs. We remark that the scale–dependence in the counterterm Lagrangian is, of course, balanced by the scale–dependence of the renormalized finite one–loop functional for observable quantities.

This method allows to control the full divergence structure of $O(q^3)$ and permits a complete renormalization of **all** Green functions: baryon form factors, photo(electro) kaon production, kaon-baryon scattering, etc. In summary, we remark that the method used has the disadvantage of leading to very lengthy and complicated expressions in the intermediate steps due to the loss of covariance. There should exist an improved method which does not share this complication. In particular, the necessary inclusion of virtual photons in the pion–nucleon (or meson–baryon) system can be treated along these lines. We hope to report on the results of such an investigation in the near future.

Acknowledgments

I am grateful to Ulf-G. Meißner for helpful discussions.

References

1. V. Bernard, N. Kaiser and Ulf-G. Meißner, Int.J.Mod.Phys. E4 (1996) 193
2. G. Ecker, Phys.Lett. B336 (1994) 508
3. G. Ecker and M. Mojzis, Phys. Lett. B365 (1996) 312
4. J. Gasser and H. Leutwyler, Ann. of Phys. 158 (1984) 142
5. J. Gasser and H. Leutwyler, Nucl. Phys. B250 (1985) 465
6. J. Gasser, M. E. Sainio and A. Svarc, Nucl.Phys B307 (1988) 779
7. E. Jenkins and A. Manohar, Phys.Lett.B255 (1991) 558
8. G. Müller and Ulf-G. Meißner, hep-ph/9610275
9. S. Weinberg, Phys.Lett. B251 (1990) 288
10. V. Bernard, N. Kaiser and Ulf-G. Meißner, Nucl. Phys. B457 (1995) 147